

contiguous complementary sequences over stretches greater than 14 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having gene-complementary stretches of 15 to 20 contiguous nucleotides, or even longer where desired.

Of course, fragments may also be obtained by other techniques such as, *e.g.*, by mechanical shearing or by restriction enzyme digestion. Small nucleic acid segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments may be obtained by application of nucleic acid reproduction technology, such as the PCR™ technology of U. S. Patents 4,683,195 and 4,683,202 (each incorporated herein by reference), by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular biology.

Accordingly, the nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of DNA fragments. Depending on the application envisioned, one will desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively stringent conditions to form the hybrids, *e.g.*, one will select relatively low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.15 M NaCl at temperatures of about 50°C to about 70°C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would be particularly suitable for isolating crystal protein-encoding DNA segments. Detection of DNA segments via hybridization is well-known to those of skill in the art, and the teachings of U. S. Patents 4,965,188 and 5,176,995 (each incorporated herein by reference) are exemplary of the methods of hybridization analyses. Teachings such as those found in the texts of Maloy *et al.*, 1994; Segal 1976; Prokop, 1991; and Kuby, 1994, are particularly relevant.

Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying template or where one seeks to isolate crystal protein-encoding sequences from related species, functional equivalents, or the like, less stringent hybridization conditions will typically be needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ conditions such as about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20°C to about 55°C. Cross-hybridizing species can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a method of choice depending on the desired results.

In certain embodiments, it will be advantageous to employ nucleic acid sequences of the present invention in combination with an appropriate means, such as a label, for determining hybridization. A wide variety of appropriate indicator means are known in the art, including fluorescent, radioactive, enzymatic or other ligands, such as avidin/biotin, which are capable of giving a detectable signal. In preferred embodiments, one will likely desire to employ a fluorescent label or an enzyme tag, such as urease, alkaline phosphatase or peroxidase, instead of radioactive or other environmental undesirable reagents. In the case of enzyme tags, colorimetric indicator substrates are known that can be employed to provide a means visible to the human eye or spectrophotometrically, to identify specific hybridization with complementary nucleic acid-containing samples.

In general, it is envisioned that the hybridization probes described herein will be useful both as reagents in solution hybridization as well as in embodiments employing a solid phase. In embodiments involving a solid phase, the test DNA (or RNA) is adsorbed or otherwise affixed to a selected matrix or surface. This fixed, single-stranded nucleic acid is then subjected to specific hybridization with selected probes under desired conditions. The selected conditions will depend on the particular circumstances based on

the particular criteria required (depending, for example, on the G+C content, type of target nucleic acid, source of nucleic acid, size of hybridization probe, *etc.*). Following washing of the hybridized surface so as to remove nonspecifically bound probe molecules, specific hybridization is detected, or even quantitated, by means of the label.

5

4.8 CHARACTERISTICS OF CRY1C* PROTEINS

The present invention provides novel polypeptides that define a whole or a portion of a *B. thuringiensis* Cry1C-R180A, Cry1C-R148A, Cry1C-R148D, Cry1C-R148L, Cry1C-R148M, Cry1C-R148G, Cry1C.563, Cry1C.499, or Cry1C.579 crystal protein.

10

In a preferred embodiment, the invention discloses and claims a purified Cry1C-R148A protein. The Cry1C-R148A protein comprises an 1189-amino acid sequence, which is given in SEQ ID NO:2.

In a second embodiment, the invention discloses and claims a purified Cry1C-R148D protein. The Cry1C-R148D protein comprises an 1189-amino acid sequence, which is given in SEQ ID NO:4.

15

In a third embodiment, the invention discloses and claims a purified Cry1C-R180A protein. The Cry1C-R180A protein comprises an 1189-amino acid sequence, which is given in SEQ ID NO:6.

In a fourth embodiment, the invention discloses and claims a purified Cry1C.563 protein. The Cry1C.563 protein comprises an 1189-amino acid sequence, which is given in SEQ ID NO:8.

20

In a fifth embodiment, the invention discloses and claims a purified Cry1C.579 protein. The Cry1C.579 protein comprises an 1189-amino acid sequence, which is given in SEQ ID NO:10.

25

In a sixth embodiment, the invention discloses and claims a purified Cry1C.499 protein. The Cry1C.499 protein comprises an 1189-amino acid sequence, which is given in SEQ ID NO:12.